

THE DEGREE OF COMMUTATIVITY OF WREATH PRODUCTS WITH INFINITE CYCLIC TOP GROUP

IKER DE LAS HERAS, BENJAMIN KLOPSCH AND ANDONI ZOZAYA

ABSTRACT. The degree of commutativity of a finite group is the probability that two uniformly and randomly chosen elements commute. This notion extends naturally to finitely generated groups G : the degree of commutativity $\text{dc}_S(G)$, with respect to a given finite generating set S , results from considering the fractions of commuting pairs of elements in increasing balls around 1_G in the Cayley graph $\mathcal{C}(G, S)$. We focus on restricted wreath products of the form $G = H \wr \langle t \rangle$, where $H \neq 1$ is finitely generated and the top group $\langle t \rangle$ is infinite cyclic. In accordance with a more general conjecture, we show that $\text{dc}_S(G) = 0$ for such groups G , regardless of the choice of S .

This extends results of Cox who considered lamplighter groups with respect to certain kinds of generating sets. We also derive a generalisation of Cox's main auxiliary result: in 'reasonably large' homomorphic images of wreath products G as above, the image of the base group has density zero, with respect to certain types of generating sets.

1. INTRODUCTION

Let G be a finitely generated group, with finite generating set S . For $n \in \mathbb{N}_0$, let $B_S(n) = B_{G,S}(n)$ denote the ball of radius n in the Cayley graph $\mathcal{C}(G, S)$ of G with respect to S . Following Antolín, Martino and Ventura [1], we define the *degree of commutativity* of G with respect to S as

$$\text{dc}_S(G) = \limsup_{n \rightarrow \infty} \frac{|\{(g, h) \in B_S(n) \times B_S(n) \mid gh = hg\}|}{|B_S(n)|^2}.$$

We remark that this notion can be viewed as a special instance of a more general concept, where the degree of commutativity is defined with respect to 'reasonable' sequences of probability measures on G , as discussed in a preliminary arXiv-version of [1] or, in more detail, by Tointon in [10].

If G is finite, the invariant $\text{dc}_S(G)$ does not depend on the particular choice of S , as the balls stabilise and $\text{dc}(G) = \text{dc}_S(G)$ simply gives the probability that two uniformly and randomly chosen elements of G commute. This situation was studied already by Erdős and Turán [4], and further results were obtained by various authors over the years; for example, see [5, 6, 7, 8]. For infinite groups G , it is generally not known whether $\text{dc}_S(G)$ is independent of the particular choice of S .

2020 *Mathematics Subject Classification.* 20P05, 20E22, 20F65, 20F69, 20F05.

Key words and phrases. Degree of commutativity, wreath products, density, word growth.

The first author acknowledges support by the Basque Government, grant POS_2021_2_0040. The third author is supported by Spanish Ministry of Science, Innovation and Universities' grant FPU17/04822. The first and third author acknowledge as well support by the Basque Government, project IT483-22, and the Spanish Government, project PID2020-117281GB-I00, partly with ERDF funds. The authors thank Heinrich-Heine-Universität Düsseldorf, where a large part of this research was carried out.

The degree of commutativity is naturally linked to the following concept of density, which is employed, for instance, in [2]. The *density* of a subset $X \subseteq G$ with respect to S is

$$\delta_S(X) = \delta_{G,S}(X) = \limsup_{n \rightarrow \infty} \frac{|X \cap B_S(n)|}{|B_S(n)|}.$$

If the group G has sub-exponential word growth, then the density function δ_S is bi-invariant; compare [2, Prop. 2.3]. Based on this fact, the following can be proved, initially for residually finite groups and then without this additional restriction, even in the more general context of suitable sequences of probability measures; see [1, Thm. 1.3] and [10, Thms. 1.6 and 1.17].

Theorem 1.1 (Antolín, Martino and Ventura [1]; Tointon [10]). *Let G be a finitely generated group of sub-exponential word growth, and let S be a finite generating set of G . Then $\text{dc}_S(G) > 0$ if and only if G is virtually abelian. Moreover, $\text{dc}_S(G)$ does not depend on the particular choice of S .*

The situation is far less clear for groups of exponential word growth. In this context, the following conjecture was raised in [1].

Conjecture 1.2 (Antolín, Martino and Ventura [1]). *Let G be a finitely generated group of exponential word growth and let S be a finite generating set of G . Then, $\text{dc}_S(G) = 0$, irrespective of the choice of S .*

In [1] the conjecture was already confirmed for non-elementary hyperbolic groups, and Valiunas [11] confirmed it for right-angled Artin groups (and more general graph products of groups) with respect to certain generating sets. Furthermore, Cox [3] showed that the conjecture holds, with respect to *selected* generating sets, for (generalised) lamplighter groups, that is for restricted standard wreath products of the form $G = F \wr \langle t \rangle$, where $F \neq 1$ is finite and $\langle t \rangle$ is an infinite cyclic group. Wreath products of such a shape are basic examples of groups of exponential word growth; in Section 2 we briefly recall the wreath product construction, here we recall that $G = A \rtimes \langle t \rangle$ with base group $A = \bigoplus_{i \in \mathbb{Z}} F^{t^i}$. In the present paper, we make a significant step forward in two directions, by confirming Conjecture 1.2 for an even wider class of restricted standard wreath products and with respect to *arbitrary* generating sets.

Theorem A. *Let $G = H \wr \langle t \rangle$ be the restricted wreath product of a finitely generated group $H \neq 1$ and an infinite cyclic group $\langle t \rangle \cong C_\infty$. Then G has degree of commutativity $\text{dc}_S(G) = 0$, for every finite generating set S of G .*

One of the key ideas in [3] is to reduce the desired conclusion $\text{dc}_S(G) = 0$, for the wreath products $G = A \rtimes \langle t \rangle$ under consideration, to the claim that the base group A has density $\delta_S(A) = 0$ in G . We proceed in a similar way and derive Theorem A from the following density result, which constitutes our main contribution.

Theorem B. *Let $G = H \wr \langle t \rangle$ be the restricted wreath product of a finitely generated group H and an infinite cyclic group $\langle t \rangle \cong C_\infty$. Then the base group $A = \bigoplus_{i \in \mathbb{Z}} H^{t^i}$ has density $\delta_S(A) = 0$ in G , for every finite generating set S of G .*

The limitation in [3] to special generating sets S of lamplighter groups G is due to the fact that the arguments used there rely on explicit minimal length expressions for elements $g \in G$ with respect to S . If one restricts to generating sets which allow control over minimal length expressions in a similar, but somewhat weaker way, it

is, in fact, possible to simplify and generalise the analysis considerably. In this way we arrive at the following improvement of the results in [3, §2.2], for homomorphic images of wreath products.

Theorem C. *Let G be a finitely generated group of exponential word growth of the form $G = A \rtimes \langle t \rangle$, where*

- (a) *the subgroup $\langle t \rangle$ is infinite cyclic;*
- (b) *the subgroup $A = \langle \bigcup \{H^{t^i} \mid i \in \mathbb{Z}\} \rangle$ is generated by the $\langle t \rangle$ -conjugates of a finitely generated subgroup H ;*
- (c) *the $\langle t \rangle$ -conjugates of this group H commute elementwise: $[H^{t^i}, H^{t^j}] = 1$ for all $i, j \in \mathbb{Z}$ with $H^{t^i} \neq H^{t^j}$.*

Suppose further that S_0 is a finite generating set for H and that the exponential growth rates of H with respect to S_0 and of G with respect to $S = S_0 \cup \{t\}$ satisfy

$$(1.1) \quad \lim_{n \rightarrow \infty} \sqrt[n]{|B_{H, S_0}(n)|} < \lim_{n \rightarrow \infty} \sqrt[n]{|B_{G, S}(n)|}.$$

Then A has density $\delta_S(A) = 0$ in G with respect to S .

For finitely generated groups G of sub-exponential word growth, the density of a subgroup of infinite index, such as A in $G = A \rtimes \langle t \rangle$ with $\langle t \rangle \cong C_\infty$, is always 0; compare [2]. Thus Theorem C has the following consequence.

Corollary 1.3. *Let $G = A \rtimes \langle t \rangle$ be a finitely generated group, where A is abelian and $\langle t \rangle \cong C_\infty$. Then A has density $\delta_S(A) = 0$ in G , with respect to any finite generating set of G that takes the form $S = S_0 \cup \{t\}$ with $S_0 \subseteq A$.*

Next we give a concrete example to illustrate that the technical condition (1.1) in Theorem C is not redundant. It is not difficult to craft more complex examples.

Example 1.4. Let $G = F \rtimes \langle t \rangle$, where $F = \langle x, y \rangle$ is the free group on two generators and $\langle t \rangle \cong C_\infty$. Then F has density $\delta_S(F) = 1/2 > 0$ in G for the ‘obvious’ generating set $S = \{x, y, t\}$.

Indeed, for every $i \in \mathbb{Z}$ we have

$$B_{G, S}(n) \cap Ft^i = \begin{cases} B_{F, \{x, y\}}(n - |i|)t^i & \text{if } n \in \mathbb{N} \text{ with } n \geq |i|, \\ \emptyset & \text{otherwise,} \end{cases}$$

and hence, for all $n \in \mathbb{N}$,

$$|B_{G, S}(n) \cap F| = |B_{F, \{x, y\}}(n)| \quad \text{and} \quad |B_{G, S}(n)| = |B_{F, \{x, y\}}(n)| + 2 \sum_{i=1}^n |B_{F, \{x, y\}}(n - i)|.$$

This yields

$$\frac{|B_{G, S}(n) \cap F|}{|B_{G, S}(n)|} = \frac{2 \cdot 3^n - 1}{2 \cdot 3^n - 1 + 2 \sum_{i=1}^n (2 \cdot 3^{n-i} - 1)} = \frac{2 \cdot 3^n - 1}{4 \cdot 3^n - 2n - 3} \rightarrow \frac{1}{2} \quad \text{as } n \rightarrow \infty.$$

We remark that in this example F and G have the same exponential growth rates:

$$\lim_{n \rightarrow \infty} \sqrt[n]{|B_{F, \{x, y\}}(n)|} = \lim_{n \rightarrow \infty} \sqrt[n]{|B_{G, S}(G)|} = 3.$$

Furthermore, the argument carries through without any obstacles with any finite generating set S_0 of F in place of $\{x, y\}$.

Finally, we record an open question that suggests itself rather naturally.

Question 1.5. Let G be a finitely generated group such that $\text{dc}_S(G) > 0$ with respect to a finite generating set S . Does it follow that there exists an abelian subgroup $A \leq G$ such that $\delta_S(A) > 0$?

For groups G of sub-exponential word growth the answer is “yes”, as one can see by an easy argument from Theorem 1.1. An affirmative answer for groups of exponential word growth could be a step towards establishing Conjecture 1.2 or provide a pathway to a possible alternative outcome. At a heuristic level, an affirmative answer to Question 1.5 would fit well with the results in [9] and [10].

Notation. Our notation is mostly standard. For elements g, h of a group G , we write $g^h = h^{-1}gh$ and $[g, h] = g^{-1}g^h$. For a finite generating set S of G , we denote by $l_S(g)$ the length of g with respect to S , i.e., the distance between g and 1 in the corresponding Cayley graph $\mathcal{C}(G, S)$.

Given $a, b \in \mathbb{R}$ and $T \subseteq \mathbb{R}$, we write $[a, b]_T = \{x \in T \mid a \leq x \leq b\}$; for instance, $[-2, \sqrt{2}]_{\mathbb{Z}} = \{-2, -1, 0, 1\}$. We repeatedly compare the limiting behaviour of real-valued functions defined on cofinite subsets of \mathbb{N}_0 which are eventually non-decreasing and take positive values. For this purpose we employ the conventional Landau symbols; specifically we write, for functions f, g of the described type,

$$f \in o(g), \text{ or } g \in \omega(f), \quad \text{if } \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0, \text{ equivalently } \lim_{n \rightarrow \infty} \frac{g(n)}{f(n)} = \infty;$$

$$f \in O(g) \quad \text{if } \limsup_{n \rightarrow \infty} \frac{f(n)}{g(n)} < \infty.$$

As customary, we use suggestive short notation such as, for instance, $f \in o(\log n)$ in place of $f \in o(g)$ for $g: \mathbb{N}_{\geq 2} \rightarrow \mathbb{R}, n \mapsto \log(n)$.

2. PRELIMINARIES

In this section, we collect preliminary and auxiliary results. Furthermore, we briefly recall the wreath product construction and establish basic notation.

2.1. Word growth. The groups G that we are concerned with have exponential word growth: for any finite generating set S of G , the *exponential growth rate*

$$(2.1) \quad \lambda_S(G) = \lim_{n \rightarrow \infty} \sqrt[n]{|B_S(n)|} = \inf \{ \sqrt[n]{|B_S(n)|} \mid n \in \mathbb{N}_0 \}$$

of G with respect to S satisfies $\lambda_S(G) > 1$. We will use the following basic estimates:

$$\lambda_S(G)^n \leq |B_S(n)| \quad \text{for all } n \in \mathbb{N}_0,$$

and, for each $\varepsilon \in \mathbb{R}_{>0}$,

$$|B_S(n)| \leq (\lambda_S(G) + \varepsilon)^n \quad \text{for all sufficiently large } n \in \mathbb{N}.$$

Furthermore we recall that $\lambda_S(G)$ has the useful alternative description

$$\lambda_S(G) = \lim_{n \rightarrow \infty} \frac{|B_S(n+1)|}{|B_S(n)|}.$$

In the proof of Theorem C, the following technical result is used.

Lemma 2.1. For each $\alpha \in [0, 1]_{\mathbb{R}}$, the sequences $\sqrt[n]{\binom{n+\lceil\alpha n\rceil}{\lceil\alpha n\rceil}}$ and $\sqrt[n]{\binom{n}{\lceil\alpha n\rceil}}$, $n \in \mathbb{N}$, converge, and furthermore

$$\lim_{\alpha \rightarrow 0^+} \left(\lim_{n \rightarrow \infty} \sqrt[n]{\binom{n+\lceil\alpha n\rceil}{\lceil\alpha n\rceil}} \right) = \lim_{\alpha \rightarrow 0^+} \left(\lim_{n \rightarrow \infty} \sqrt[n]{\binom{n}{\lceil\alpha n\rceil}} \right) = 1.$$

Consequently, if $f: \mathbb{N} \rightarrow \mathbb{R}_{>0}$ satisfies $f \in o(n)$, then the sequence $\binom{n+\lceil f(n)\rceil}{\lceil f(n)\rceil}$, $n \in \mathbb{N}$, grows sub-exponentially in n , viz. $\sqrt[n]{\binom{n+\lceil f(n)\rceil}{\lceil f(n)\rceil}} \rightarrow 1$ as $n \rightarrow \infty$.

Proof. For each $\alpha \in [0, 1]_{\mathbb{R}}$, Stirling's approximation for factorials yields

$$\begin{aligned} \binom{n}{\lceil\alpha n\rceil} &\sim \frac{\sqrt{2\pi n}(n/e)^n}{\sqrt{2\pi\lceil\alpha n\rceil}(\lceil\alpha n\rceil/e)^{\lceil\alpha n\rceil} \sqrt{2\pi(n-\lceil\alpha n\rceil)}((n-\lceil\alpha n\rceil)/e)^{(n-\lceil\alpha n\rceil)}} \\ &= \frac{\sqrt{n}}{\sqrt{2\pi\lceil\alpha n\rceil\lceil n-\alpha n\rceil}} \frac{n^n}{\lceil\alpha n\rceil^{\lceil\alpha n\rceil} \lceil n-\alpha n\rceil^{\lceil n-\alpha n\rceil}}, \quad \text{as } n \rightarrow \infty, \end{aligned}$$

i.e., the ratio of the left-hand term to the right-hand term tends to 1 as n tends to infinity. Moreover, for all sufficiently large $n \in \mathbb{N}$,

$$\begin{aligned} \frac{n^n}{\lceil\alpha n\rceil^{\lceil\alpha n\rceil} \lceil n-\alpha n\rceil^{\lceil n-\alpha n\rceil}} &\leq \frac{n^n}{(\alpha n)^{\alpha n} (n-\alpha n-1)^{n-\alpha n-1}} \\ &= n \left(\frac{1}{\alpha^\alpha (1-\alpha-1/n)^{(1-\alpha-1/n)}} \right)^n. \end{aligned}$$

This shows that

$$\lim_{n \rightarrow \infty} \sqrt[n]{\binom{n}{\lceil\alpha n\rceil}} \leq \frac{1}{\alpha^\alpha (1-\alpha)^{1-\alpha}}.$$

A similar computation yields

$$\lim_{n \rightarrow \infty} \sqrt[n]{\binom{n+\lceil\alpha n\rceil}{\lceil\alpha n\rceil}} \leq \frac{(1+\alpha)^{1+\alpha}}{\alpha^\alpha}.$$

Since $\lim_{\alpha \rightarrow 0^+} \alpha^\alpha = 1$, we conclude that

$$\lim_{\alpha \rightarrow 0^+} \left(\lim_{n \rightarrow \infty} \sqrt[n]{\binom{n+\lceil\alpha n\rceil}{\lceil\alpha n\rceil}} \right) = \lim_{\alpha \rightarrow 0^+} \left(\lim_{n \rightarrow \infty} \sqrt[n]{\binom{n}{\lceil\alpha n\rceil}} \right) = 1. \quad \square$$

2.2. Wreath products. We recall that a group $G = H \wr K$ is the restricted standard wreath product of two subgroups H and K , if it decomposes as a semidirect product $G = A \rtimes K$, where the normal closure of H is the direct sum $A = \bigoplus_{k \in K} H^k$ of the various conjugates of H by elements of K ; the groups A and K are referred to as the *base group* and the *top group* of the wreath product G , respectively. Since we do not consider complete standard wreath products or more general types of permutational wreath products, we shall drop the terms ‘‘restricted’’ and ‘‘standard’’ from now on.

Throughout the rest of this section, let

$$(2.2) \quad G = H \wr \langle t \rangle = A \rtimes \langle t \rangle \quad \text{with base group} \quad A = \bigoplus_{i \in \mathbb{Z}} H^{t^i}$$

be the wreath product of a finitely generated subgroup H and an infinite cyclic subgroup $\langle t \rangle \cong C_\infty$. Every element $g \in G$ can be written uniquely in the form

$$g = \tilde{g} t^{\rho(g)}, \quad \text{where } \rho(g) \in \mathbb{Z} \text{ and } \tilde{g} = \prod_{i \in \mathbb{Z}} (g_i)^{t^i} \in A \text{ with ‘coordinates’ } g_i \in H.$$

The support of the product decomposition of \tilde{g} is finite and we write

$$\text{supp}(g) = \{i \in \mathbb{Z} \mid g_i \neq 1\}.$$

Furthermore, it is convenient to fix a finite symmetric generating set S of G ; thus $G = \langle S \rangle$, $1 \in S$, and $g \in S$ implies $g^{-1} \in S$. We put $d = |S|$ and fix an ordering of the elements of S :

$$(2.3) \quad S = \{s_1, \dots, s_d\}, \quad \text{with } s_j = \tilde{s}_j t^{\rho(s_j)} \text{ for } j \in [1, d]_{\mathbb{Z}},$$

where $\tilde{s}_1, \dots, \tilde{s}_d \in A$. We write $r_S = \max \{\rho(s_j) \mid j \in [1, d]_{\mathbb{Z}}\}$.

Definition 2.2. An S -expression of an element $g \in G$ is (induced by) a word $W = \prod_{k=1}^l X_{\iota(k)}$ in the free semigroup $\langle X_1, \dots, X_d \rangle$ such that

$$(2.4) \quad g = W(s_1, \dots, s_d) = \prod_{k=1}^l s_{\iota(k)};$$

here W determines $\iota = \iota_W: [1, l]_{\mathbb{Z}} \rightarrow [1, d]_{\mathbb{Z}}$ and vice versa. We observe that in this situation

$$(2.5) \quad g = \tilde{g} t^{-\sigma(l)} \quad \text{with} \quad \tilde{g} = \prod_{k=1}^l \widetilde{s_{\iota(k)}} t^{\sigma(k-1)},$$

where $\sigma = \sigma_{S,W}$ is short for the negative¹ cumulative exponent function

$$\sigma_{S,W}: [0, l]_{\mathbb{Z}} \rightarrow \mathbb{Z}, \quad k \mapsto - \sum_{j=1}^k \rho(s_{\iota(j)}).$$

We define the *itinerary* of g associated to the S -expression (2.4) as the pair

$$\text{It}(S, W) = (\iota_W, \sigma_{S,W}),$$

and we say that $\text{It}(S, W)$ has length l , viz. the length of the word W . The terminology refers to (2.5), which indicates how g can be built stepwise from the ‘sequences’ ι_W and $\sigma_{S,W}$; see Example 2.4 below. Furthermore we call

$$\text{maxi}(\text{It}(S, W)) = \max(\sigma_{S,W}) \quad \text{and} \quad \text{mini}(\text{It}(S, W)) = \min(\sigma_{S,W})$$

the *maximal* and *minimal itinerary points* of $\text{It}(S, W)$.

Later we fix a representative function $\mathcal{W}: G \rightarrow \langle X_1, \dots, X_d \rangle$, $g \mapsto W_g$ which yields for each element of G an S -expression of shortest possible length. In that situation we suppress the reference to S and refer to

$$\text{It}_{\mathcal{W}}(g) = \text{It}(S, W_g), \quad \text{maxi}_{\mathcal{W}}(g) = \text{maxi}(\text{It}_{\mathcal{W}}(g)), \quad \text{mini}_{\mathcal{W}}(g) = \text{mini}(\text{It}_{\mathcal{W}}(g))$$

as the \mathcal{W} -itinerary, the *maximal \mathcal{W} -itinerary point* and the *minimal \mathcal{W} -itinerary point* of any given element g .

Remark 2.3. (1) If $I = (\iota, \sigma)$ is the itinerary of an element $g \in G$ associated to some S -expression of length l , then g is uniquely determined by I via (2.5). Hence, we call g the element corresponding to the itinerary I .

¹At this stage the sign change is a price we pay for not introducing notation for left-conjugation; Example 2.4 illustrates that σ plays a convenient role in the concept of itinerary.

(2) Suppose that $I_1 = (\iota_1, \sigma_1)$ and $I_2 = (\iota_2, \sigma_2)$ are itineraries, of lengths l_1 and l_2 , associated to S -expressions W_1, W_2 for elements $g_1, g_2 \in G$. Clearly, $W = W_1 W_2$ is an S -expression for $g = g_1 g_2$; the associated itinerary

$$I = \text{It}(S, W) = (\iota, \sigma)$$

has length $l = l_1 + l_2$ and its components are given by

$$\iota(k) = \begin{cases} \iota_1(k) & \text{if } k \in [1, l_1]_{\mathbb{Z}}, \\ \iota_2(k - l_1) & \text{if } k \in [l_1 + 1, l]_{\mathbb{Z}}, \end{cases}$$

$$\sigma(k) = \begin{cases} \sigma_1(k) & \text{if } k \in [0, l_1]_{\mathbb{Z}}, \\ \sigma_1(l_1) + \sigma_2(k - l_1) & \text{if } k \in [l_1 + 1, l]_{\mathbb{Z}}. \end{cases}$$

We refer to I as the *product itinerary* and write $I = I_1 * I_2$.

Conversely, if I is the itinerary of some element $g \in G$ associated to some S -expression of length l and if $l_1 \in [0, l]_{\mathbb{Z}}$, there is a unique decomposition $I = I_1 * I_2$ for itineraries I_1 of length l_1 and I_2 of length $l_2 = l - l_1$; the corresponding elements $g_1, g_2 \in G$ satisfy $g = g_1 g_2$.

(3) For a representative function $\mathcal{W}: G \rightarrow \langle X_1, \dots, X_d \rangle$, $g \mapsto W_g$, as discussed in Definition 2.2, it is typically not the case that $W_{gh} = W_g W_h$ for $g, h \in G$ and, consequently, neither $\text{It}_{\mathcal{W}}(gh) = \text{It}_{\mathcal{W}}(g) * \text{It}_{\mathcal{W}}(h)$ holds.

Example 2.4. A typical example of the wreath products that we consider is the lamplighter group

$$G = \langle a, t \mid a^2 = 1, [a, a^{t^i}] = 1 \text{ for } i \in \mathbb{Z} \rangle = \bigoplus_{i \in \mathbb{Z}} \langle a_i \rangle \rtimes \langle t \rangle \cong C_2 \wr C_{\infty},$$

where $a_i = a^{t^i}$ for each $i \in \mathbb{Z}$. We consider the finite symmetric generating set

$$S = \{s_1, \dots, s_6\} \quad \text{with} \quad s_1 = a_4 t^{-3}, s_2 = t^{-2}, s_3 = 1, s_4 = a_0, s_5 = t^2, s_6 = a_1 t^3.$$

Let $g = \tilde{g} t^3$ be such that $g_{|i} = a$ for $i \in \{-1, 1, 2, 6\}$ and $g_{|i} = 1$ otherwise. Then we have $\rho(g) = 3$, $\text{supp}(g) = \{-1, 1, 2, 6\}$, and

$$(2.6) \quad g = t^{-2} \cdot a_0 \cdot a_4 t^{-3} \cdot t^2 \cdot t^2 \cdot a_0 \cdot t^2 \cdot a_0 \cdot t^2 = s_2 s_4 s_1 s_5 s_5 s_4 s_5 s_4 s_5$$

is an S -expression for g of length 9, based on $W = X_2 X_4 X_1 X_5 X_5 X_4 X_5 X_4 X_5$. The itinerary $I = \text{It}(S, W)$ associated to this S -expression for g is

$$(2.7) \quad I = (\iota, \sigma) = ((2, 4, 1, 5, 5, 4, 5, 4, 5), (0, 2, 2, 5, 3, 1, 1, -1, -1, -3)),$$

where we have written the maps ι and σ in sequence notation. Furthermore, we see that $\text{maxi}(I) = 5$ and $\text{mini}(I) = -3$. Figure 1 gives a pictorial description of part of the information contained in I .




FIGURE 1. An illustration of the itinerary (2.7) associated to the S -expression (2.6).

Lemma 2.5. *Let $G = H \wr \langle t \rangle$ be a wreath product as in (2.2), with generating set S as in (2.3). Then the following hold.*

(i) Put

$$C = C(S) = 1 + \max \{|i| \mid i \in \text{supp}(s) \text{ for some } s \in S\} \in \mathbb{N}.$$

Then for every $g \in G$ with itinerary I ,

$$\text{mini}(I) - C < \min(\text{supp}(g)) \quad \text{and} \quad \max(\text{supp}(g)) < \text{maxi}(I) + C.$$

(ii) Let $u \in H$ and put

$$D = D(S, u) = l_S(u) + 2 \max \{l_S(t^j) \mid -r_S \leq j \leq r_S\} \in \mathbb{N}.$$

Let $g \in G$ with itinerary I , associated to an S -expression of length $l_S(g)$. Then, for every $j \in \mathbb{Z}$ with $\text{mini}(I) - r_S \leq j \leq \text{maxi}(I) + r_S$, the elements $h = gu^{t^{j+\rho(g)}}$, $\tilde{h} = u^{t^j}g \in G$ satisfy $\rho(h) = \rho(\tilde{h}) = \rho(g)$ and the ‘coordinates’ of h, \tilde{h} are given by

$$h_{|i} = \begin{cases} g_{|i} & \text{if } i \neq j, \\ g_{|j} u & \text{if } i = j, \end{cases} \quad \tilde{h}_{|i} = \begin{cases} g_{|i} & \text{if } i \neq j, \\ u g_{|j} & \text{if } i = j \end{cases} \quad \text{for } i \in \mathbb{Z}.$$

Furthermore,

$$l_S(h) \leq l_S(g) + D \quad \text{and} \quad l_S(\tilde{h}) \leq l_S(g) + D.$$

Proof. We write $I = (\iota, \sigma)$, and l denotes the length of I .

(i) From (2.5) it follows that

$$\begin{aligned} \text{supp}(g) &\subseteq \bigcup_{1 \leq k \leq l} (\sigma(k-1) + \text{supp}(s_{\iota(k)})) \\ &\subseteq \bigcup_{1 \leq k \leq l} [\sigma(k-1) - C + 1, \sigma(k-1) + C - 1]_{\mathbb{Z}}; \end{aligned}$$

from this inclusion the two inequalities follow readily.

(ii) The first assertions are very easy to verify. We justify the upper bound for $l_S(h)$; the bound for $l_S(\tilde{h})$ follows similarly.

Since $\text{mini}(I) - r_S \leq j \leq \text{maxi}(I) + r_S$ and since itineraries proceed, in the second coordinate, by steps of length at most r_S , there exists $k \in [0, l]_{\mathbb{Z}}$ such that $|j - \sigma(k)| \leq r_S$; we choose k maximal with this property. Next we decompose the itinerary I as the product $I = I_1 * I_2$ of itineraries I_1 of length $l_1 = k$ and I_2 of length $l_2 = l - k$; compare Remark 2.3. Choose an itinerary I_3 , associated to an S -expression for $t^{-j+\sigma(k)} u t^{j-\sigma(k)}$ of length $l_3 \leq l_S(u) + 2l_S(t^{j-\sigma(k)})$. Then $I_1 * I_3 * I_2$ is an itinerary for h , associated to an S -expression of length $l_1 + l_2 + l_3$; this uses that k was chosen maximal. We conclude that

$$l_S(h) \leq l + l_3 \leq l_S(g) + l_S(u) + 2l_S(t^{j-\sigma(k)}) \leq l_S(g) + D. \quad \square$$

3. PROOFS OF THEOREMS A AND B

First we explain how Theorem A follows from Theorem B. The first ingredient is the following general lemma.

Lemma 3.1 (Antolín, Martino and Ventura [1, Lem. 3.1]). *Let $G = \langle S \rangle$ be a group, with finite generating set S . Suppose that there exists a subset $X \subseteq G$ satisfying*

- (a) $\delta_S(X) = 0$;
- (b) $\sup \left\{ \frac{|C_G(g) \cap B_S(n)|}{|B_S(n)|} \mid g \in G \setminus X \right\} \rightarrow 0$ as $n \rightarrow \infty$.

Then G has degree of commutativity $\text{dc}_S(G) = 0$.

The second ingredient comes from [3, §2.1], where Cox shows the following. If $G = H \wr \langle t \rangle$ is the wreath product of a finitely generated group $H \neq 1$ and an infinite cyclic group $\langle t \rangle$, with base group A , and if S is any finite generating set for G , then

$$\sup \left\{ \frac{|C_G(g) \cap B_S(n)|}{|B_S(n)|} \mid g \in G \setminus A \right\} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Thus, Theorem B implies Theorem A.

It remains to establish Theorem B. Throughout the rest of this section, let

$$G = H \wr \langle t \rangle = A \rtimes \langle t \rangle \quad \text{with base group} \quad A = \bigoplus_{i \in \mathbb{Z}} H^{t^i}$$

be the wreath product of a finitely generated subgroup H and an infinite cyclic subgroup $\langle t \rangle$, just as in (2.2). The exceptional case $H = 1$ poses no obstacle, hence we assume $H \neq 1$. We fix a finite symmetric generating set $S = \{s_1, \dots, s_d\}$ for G and employ the notation established around (2.3). Finally, we recall that G has exponential word growth and we write

$$\lambda = \lambda_S(G) > 1$$

for the exponential growth rate of G with respect to S ; compare (2.1).

We start by showing that the subset of A consisting of all elements with suitably bounded support is negligible in the computation of the density of A .

Lemma 3.2. *Fix a representative function \mathcal{W} which yields for each element of G an S -expression of shortest possible length. Let $q: \mathbb{N} \rightarrow \mathbb{R}_{\geq 1}$ be a non-decreasing unbounded function such that $q \in o(\log n)$. For $n \in \mathbb{N}$, we consider*

$$R_q(n) = R_{\mathcal{W},q}(n) = \{g \in A \cap B_S(n) \mid \max_{\mathcal{W}}(g) - \min_{\mathcal{W}}(g) \leq q(n)\}.$$

Then

$$\lim_{n \rightarrow \infty} \frac{|R_q(n)|}{|B_S(n)|} = 0.$$

Proof. For $i \in \mathbb{Z}$, we write $H_i = H^{t^i}$. Using the notation established in Section 2.2, we accumulate the ‘coordinates’ of elements of S in

$$S_0 = \{s_{|i} \mid s \in S, i \in \mathbb{Z}\} = \{(s_j)_{|i} \mid 1 \leq j \leq d \text{ and } i \in \mathbb{Z}\} \subseteq H = H_0,$$

and we set $S_i = S_0^{t^i} \subseteq H_i$ for $i \in \mathbb{Z}$. Then S_i is a finite symmetric generating set of H_i for each $i \in \mathbb{Z}$. Moreover, we have $|B_{H_i, S_i}(n)| = |B_{H, S_0}(n)|$ for all $n \in \mathbb{N}$; consequently,

$$\lambda_{S_0}(H) = \lambda_{S_i}(H_i) \quad \text{for all } i \in \mathbb{Z}.$$

Let $C = C(S) \in \mathbb{N}$ be as is in Lemma 2.5(i), and choose a non-decreasing unbounded function $f: \mathbb{N} \rightarrow \mathbb{R}_{>0}$ such that

$$f \in o(n/q(n)) \quad \text{and} \quad f \in \omega((\lambda + 1)^{m(n)}) \quad \text{for } m(n) = (3q(n) + 4C)l_S(n);$$

for instance, we could take $f = f_\alpha$ for any real parameter α with $0 < \alpha < 1$, where $f_\alpha(n) = \max \{k^\alpha/q(k) \mid k \in [1, n]_{\mathbb{Z}}\}$ for $n \in \mathbb{N}$. We consider

$$R_q^f(n) = R_{\mathcal{W},q}^f(n) = \{g \in R_q(n) \mid g_{|i} \in B_{H, S_0}(f(n)) \text{ for all } i \in \mathbb{Z}\}.$$

Choose $C' \in \mathbb{N}$ such that $\lambda^{C'} > \lambda_{S_0}(H)$. Then we have, for all sufficiently large $n \in \mathbb{N}$,

$$|R_q^f(n)| \leq |B_{H, S_0}(f(n))|^{2q(n)+2C} \leq \lambda^{2C'q(n)f(n)+2C' Cf(n)} \leq \lambda^{4C' C q(n)f(n)}.$$

From $f \in o(n/q(n))$ we obtain

$$4C' C q(n)f(n) - n \rightarrow -\infty \quad \text{as } n \rightarrow \infty$$

and hence

$$\frac{|R_q^f(n)|}{|B_S(n)|} \leq \lambda^{4C' C q(n) f(n) - n} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

It remains to show that

$$\frac{|R_q(n) \setminus R_q^f(n)|}{|B_S(n)|} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

For every $g \in R_q(n) \setminus R_q^f(n)$ we pick $i(g) \in \mathbb{Z}$ with

$$\text{mini}_{\mathcal{W}}(g) - C < i(g) < \text{maxi}_{\mathcal{W}}(g) + C \quad \text{and} \quad g_{|i(g)} \notin B_{S_0}(f(n)).$$

Let $I = (\iota, \sigma)$, viz. $I_g = (\iota_g, \sigma_g)$, denote the \mathcal{W} -itinerary of g . Then

$$g_{|i(g)} = \prod_{k=1}^{l_S(g)} (s_{\iota(k)})_{|i(g) - \sigma(k-1)}.$$

By successively cancelling sub-products of adjacent factors that evaluate to 1 and have maximal length with this property (in an orderly fashion, from left to right, say), we arrive at a ‘reduced’ product expression

$$(3.1) \quad g_{|i(g)} = \prod_{j=1}^{\ell} (s_{\iota(\kappa(j))})_{|i(g) - \sigma(\kappa(j)-1)},$$

for some $\ell = \ell_g \in [1, l_S(g)]_{\mathbb{Z}}$ and an increasing function $\kappa = \kappa_g: [1, \ell]_{\mathbb{Z}} \rightarrow [1, l_S(g)]_{\mathbb{Z}}$ that picks out a subsequence of factors. In particular, this means that, for $j_1, j_2 \in [1, \ell]_{\mathbb{Z}}$ with $j_1 < j_2$,

$$(3.2) \quad \begin{aligned} \prod_{k=\kappa(j_1)+1}^{\kappa(j_2)} (s_{\iota(k)})_{|i(g) - \sigma(k-1)} &= \prod_{j=j_1+1}^{j_2} \prod_{k=\kappa(j-1)+1}^{\kappa(j)} (s_{\iota(k)})_{|i(g) - \sigma(k-1)} \\ &= \prod_{j=j_1+1}^{j_2} (s_{\iota(\kappa(j))})_{|i(g) - \sigma(\kappa(j)-1)} \neq 1, \end{aligned}$$

and moreover we have $l_S(g) \geq \ell \geq l_{S_0}(g_{|i(g)}) \geq f(n)$.

For each choice of $j \in [1, \ell]_{\mathbb{Z}}$ we decompose the itinerary I for g into a product $I = I_{j,1} * I_{j,2}$ of itineraries of length $\kappa(j)$ and $l_S(g) - \kappa(j)$; compare Remark 2.3. Then $g = h_{j,1} h_{j,2}$, where $h_{j,1}, h_{j,2}$ denote the elements of G corresponding to $I_{j,1}, I_{j,2}$. From $g \in R_q(n)$ it follows that $\text{maxi}(I_{j,1}) - \text{mini}(I_{j,1})$ and $\text{maxi}(I_{j,2}) - \text{mini}(I_{j,2})$ are bounded by $q(n)$; in particular, $\rho(h_{j,1}) \in [-q(n), q(n)]_{\mathbb{Z}}$.

We define

$$\dot{g}(j) = h_{j,1} t^{-3q(n)-4C} h_{j,2} \in B_S(n + m(n)),$$

where C denotes again the constant introduced in Lemma 2.5(i) and $m(n)$ is as defined above.

We prove below:

- (i) for each $j \in [1, \ell]_{\mathbb{Z}}$ the original element g can be recovered from $\dot{g}(j)$;
- (ii) the elements $\dot{g}(1), \dots, \dot{g}(\ell)$ are pairwise distinct.

Based on these assertions, the proof concludes as follows. We define a map

$$\begin{aligned} F_n: R_q(n) \setminus R_q^f(n) &\rightarrow \mathcal{P}(B_S(n + m(n))) \\ g &\mapsto \{\dot{g}(j) \mid 1 \leq j \leq \ell_g\}. \end{aligned}$$

From (i) we deduce that $F_n(g_1) \cap F_n(g_2) = \emptyset$ for all $g_1, g_2 \in R_q(n) \setminus R_q^f(n)$ with $g_1 \neq g_2$. In addition, from $\ell_g \geq f(n)$ and (ii) we deduce that $|F_n(g)| \geq f(n)$ for all $g \in R_q(n) \setminus R_q^f(n)$. This yields

$$|B_S(n + m(n))| \geq f(n) |R_q(n) \setminus R_q^f(n)|,$$

and hence,

$$\begin{aligned} \frac{|R_q(n) \setminus R_q^f(n)|}{|B_S(n)|} &\leq \frac{|B_S(n + m(n))|}{f(n) |B_S(n)|} \leq \frac{|B_S(m(n))|}{f(n)} \\ &\leq \frac{(\lambda + 1)^{m(n)}}{f(n)} \rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned}$$

Thus, it suffices to justify (i) and (ii), for fixed $g \in G$.

(i) Let $j \in [1, \ell]_{\mathbb{Z}}$, and write $\mathcal{H}_1 = \text{supp}(h_{j,1})$, $\mathcal{H}_2 = \text{supp}(h_{j,2})$. Lemma 2.5(i) implies that the sets \mathcal{H}_1 and $\mathcal{H}_2 - \rho(h_{j,1}) = \text{supp}(t^{\rho(h_{j,1})} h_{j,2})$ lie wholly within the interval $[-q(n) - C, q(n) + C]_{\mathbb{Z}}$, hence

$$(3.3) \quad \text{supp}(\dot{g}(j)) = \mathcal{H}_1 \cup (\mathcal{H}_2 - \rho(h_{j,1}) + 3q(n) + 4C)$$

with a gap

$$\begin{aligned} \tau &= \underbrace{\min(\mathcal{H}_2 - \rho(h_{j,1}) + 3q(n) + 4C)}_{\geq -q(n) - C + 3q(n) + 4C = 2q(n) + 3C} - \underbrace{\max(\mathcal{H}_1)}_{\leq q(n) + C} \geq q(n) + 2C, \end{aligned}$$

subject to the standard conventions $\min \emptyset = +\infty$ and $\max \emptyset = -\infty$ in special circumstances; see Figure 2 for a pictorial illustration.




FIGURE 2. An illustration of the factorisation $\dot{g}(j) = h_{j,1} t^{-3q(n)-4C} h_{j,2}$.

In contrast, gaps between two elements in \mathcal{H}_1 or two elements in \mathcal{H}_2 are strictly less than $q(n) + 2C$. Consequently, we can identify the two components in (3.3) and thus \mathcal{H}_1 and $\mathcal{H}_2 - \rho(h_{j,1})$, without any prior knowledge of j or $h_{j,1}, h_{j,2}$. Therefore, for each $i \in \mathbb{Z}$ the i th coordinate of g satisfies

$$g_i = \begin{cases} \dot{g}(j)|_i \dot{g}(j)|_{i+3q(n)+4C} & \text{if } i \in [-q(n) - C, q(n) + C], \\ 1 & \text{otherwise,} \end{cases}$$

and hence g can be recovered from $\dot{g}(j)$.

(ii) For $j_1, j_2 \in [1, \ell]_{\mathbb{Z}}$ with $j_1 < j_2$ we conclude from our choice of the ‘reduced’ product expression (3.1) and its consequence (3.2) that

$$\dot{g}(j_1)|_{i(g)} = \prod_{k=1}^{\kappa(j_1)} (s_{\iota(k)})|_{i(g)-\sigma(k-1)} \neq \prod_{k=1}^{\kappa(j_2)} (s_{\iota(k)})|_{i(g)-\sigma(k-1)} = \dot{g}(j_2)|_{i(g)},$$

and hence $\dot{g}(j_1) \neq \dot{g}(j_2)$. □

Remark 3.3. Lemma 3.2 can be established much more easily under the extra assumption that H has sub-exponential word growth. Indeed, in this case, one can prove that

$$\lim_{n \rightarrow \infty} \frac{|R_q(n)|}{|B_S(n)|} = 0$$

for any non-decreasing unbounded function $q: \mathbb{N} \rightarrow \mathbb{R}_{>1}$ such that $q \in o(n)$; the proof is similar to the one of Lemma 4.1 below.

If we assume that H is finite, it is easy to see that there exists $\alpha \in \mathbb{R}_{>0}$ such that

$$\lim_{n \rightarrow \infty} \frac{|R_q(n)|}{|B_S(n)|} = 0 \quad \text{for } q: \mathbb{N} \rightarrow \mathbb{R}_{>1}, n \mapsto 1 + \alpha n.$$

Next we establish Theorem B, using ideas that are similar to those in the proof of Lemma 3.2.

Proof of Theorem B. Fix a representative function \mathcal{W} which yields for each element of G an S -expression of shortest possible length. Let $q: \mathbb{N} \rightarrow \mathbb{R}_{\geq 1}$ be a non-decreasing unbounded function such that $q \in o(\log n)$. We make use of the decomposition

$$(3.4) \quad A \cap B_S(n) = R_q(n) \cup R_q^b(n), \quad \text{for } n \in \mathbb{N},$$

where $R_q(n) = R_{\mathcal{W},q}(n)$ is defined as in Lemma 3.2 and $R_q^b(n) = R_{\mathcal{W},q}^b(n)$ denotes the corresponding complement. Indeed, our aim is to produce a constant $D' \in \mathbb{N}$ such that

$$(3.5) \quad |B_S(n + D')| > \frac{q(n)}{2} |R_q^b(n)| \quad \text{for } n \in \mathbb{N}.$$

This yields

$$\frac{|R_q^b(n)|}{|B_S(n)|} < \frac{2|B_S(n + D')|}{q(n)|B_S(n)} \leq \frac{2|B_S(D')|}{q(n)} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Together with Lemma 3.2 we deduce from (3.4) that

$$\delta_S(A) = \lim_{n \rightarrow \infty} \frac{|A \cap B_S(n)|}{|B_S(n)|} = 0.$$

For later use, we fix an element $u \in H \setminus \{1\}$. Let $n \in \mathbb{N}$ and consider $g \in R_q^b(n)$ with \mathcal{W} -itinerary $I = (\iota, \sigma)$, viz. $I_g = (\iota_g, \sigma_g)$. We put

$$\sigma^+ = \sigma_g^+ = \max_{\mathcal{W}}(g) \quad \text{and} \quad \sigma^- = \sigma_g^- = \min_{\mathcal{W}}(g).$$

For the time being, we suppose that

$$\begin{aligned} k^+ &= k_{\mathcal{W},g}^+ = \min\{k \mid 0 \leq k \leq l_S(g) \text{ and } \sigma(k) = \sigma^+\}, \\ k^- &= k_{\mathcal{W},g}^- = \min\{k \mid 0 \leq k \leq l_S(g) \text{ and } \sigma(k) = \sigma^-\} \end{aligned}$$

satisfy $k^+ \leq k^-$; if $k^- < k^+$, a similar argument can be applied as we shall see in the course of the proof. We decompose the itinerary for g as $I = I_1 * I_2 * I_3$, where I_1, I_2, I_3 have lengths $k^+, k^- - k^+, l_S(g) - k^-$; compare Remark 2.3. If $x = x_{\mathcal{W},g}, y = y_{\mathcal{W},g}, z = z_{\mathcal{W},g}$ denote the elements corresponding to I_1, I_2, I_3 then $g = xyz$; observe that the lengths of I_1, I_2, I_3 are automatically minimal, i.e, equal to $l_S(x), l_S(y), l_S(z)$. All this is illustrated schematically in Figure 3.




FIGURE 3. A schematic illustration of the decomposition $g = xyz$. Observe that I_1 , associated to x , ‘starts’ at 0 and ‘ends’ at σ^+ , the shifted I_2 , associated to y , ‘starts’ at σ^+ and ‘ends’ at σ^- , and the shifted I_3 , associated to z , ‘starts’ at σ^- and ‘ends’ at 0.

Next, we put to use the element $u \in H \setminus \{1\}$ that was fixed and define, for any given $J \subseteq \mathbb{Z}$, perturbations

$$\dot{x}(J) = \dot{x}_{\mathcal{W},g}(J, u), \quad \dot{y}(J) = \dot{y}_{\mathcal{W},g}(J, u), \quad \dot{z}(J) = \dot{z}_{\mathcal{W},g}(J, u)$$

of the elements x, y, z that are specified by

$$(3.6) \quad \rho(\dot{x}(J)) = \rho(x) = -\sigma^+, \quad \rho(\dot{y}(J)) = \rho(y) = -\sigma^- + \sigma^+, \quad \rho(\dot{z}(J)) = \rho(z) = \sigma^-$$

and

$$(3.7) \quad \begin{aligned} \dot{x}(J)|_i &= \begin{cases} x|_i u & \text{for } i \in J_{\geq 0}, \\ x|_i & \text{otherwise,} \end{cases} \\ \dot{y}(J)|_i &= \begin{cases} u^{-1} y|_i & \text{for } i \in \mathbb{Z} \text{ such that } i + \sigma^+ \in J_{\geq 0}, \\ y|_i u^{-1} & \text{for } i \in \mathbb{Z} \text{ such that } i + \sigma^+ \in J_{< 0}, \\ y|_i & \text{otherwise,} \end{cases} \\ \dot{z}(J)|_i &= \begin{cases} u z|_i & \text{for } i \in \mathbb{Z} \text{ such that } i + \sigma^- \in J_{< 0}, \\ z|_i & \text{otherwise.} \end{cases} \end{aligned}$$

We observe that

$$(3.8) \quad g = \dot{x}(J) \dot{y}(J) \dot{z}(J).$$

Let $C = C(S) \in \mathbb{N}$ be as is in Lemma 2.5(i), and define

$$\dot{g}(J) = \dot{x}(J) t^{-2C} \dot{y}(J)^{-1} t^{-2C} \dot{z}(J);$$

compare Figure 4.




FIGURE 4. A schematic illustration of $\dot{g}(J)$, ‘starting’ at 0 and ‘ending’ at $-\rho(\dot{g}(J))$.

Suppose now that $J \subseteq [\sigma^-, \sigma^+]_{\mathbb{Z}}$ with $|J| = 2$. Let $D = D(S, u) \in \mathbb{N}$ be as is in Lemma 2.5(ii). Since

$$\begin{aligned} J_{\geq 0} &\subseteq [0, \sigma^+]_{\mathbb{Z}} \subseteq [\text{mini}(I_1), \text{maxi}(I_1)]_{\mathbb{Z}}, \\ J - \sigma^+ &\subseteq [\sigma^- - \sigma^+, 0]_{\mathbb{Z}} = [\text{mini}(I_2), \text{maxi}(I_2)]_{\mathbb{Z}}, \\ J_{< 0} - \sigma^- &\subseteq [0, -\sigma^-]_{\mathbb{Z}} \subseteq [\text{mini}(I_3), \text{maxi}(I_3)]_{\mathbb{Z}} \end{aligned}$$

we can apply Lemma 2.5(ii), if necessary twice, to deduce that

$$l_S(\dot{x}(J)) \leq l_S(x) + 2D, \quad l_S(\dot{y}(J)) \leq l_S(y) + 2D, \quad l_S(\dot{z}(J)) \leq l_S(z) + 2D.$$

Since $l_S(x) + l_S(y) + l_S(z) \leq l_S(g)$, this gives

$$l_S(\dot{g}(J)) \leq l_S(g) + D' \leq n + D' \quad \text{for } D' = 6D + 2l_S(t^{2C})$$

and hence $\dot{g}(J) \in B_S(n + D')$.

Observe that

$$\dot{g}(J) \in At^{\rho(\dot{g}(J))}, \quad \text{where } \rho(\dot{g}(J)) = 2(\sigma_g^+ - \sigma_g^-) + 4C \geq 4.$$

Up to now we assumed that $k^+ \leq k^-$; if instead $k^- < k^+$, a similar construction at this stage yields elements

$$\dot{g}(J) \in At^{\rho(\dot{g}(J))}, \quad \text{where } \rho(\dot{g}(J)) = 2(\sigma_g^- - \sigma_g^+) - 4C \leq -4;$$

in particular, there is no overlap between elements $\dot{g}(J)$ arising from these two different cases.

The set $R_q^b(n)$ decomposes into a disjoint union of subsets

$$R_{q,\ell}^b(n) = \{g \in A \cap B_S(n) \mid \sigma_g^+ - \sigma_g^- = \ell\}, \quad \ell > q(n),$$

and the map

$$\begin{aligned} F_n: R_q^b(n) &\rightarrow \mathcal{P}(B_S(n + D')), \\ g &\mapsto \{\dot{g}(J) \mid J \subseteq [\sigma_g^-, \sigma_g^+]_{\mathbb{Z}} \text{ with } |J| = 2\} \end{aligned}$$

restricts, for each $\ell \in \mathbb{N}$ with $\ell > q(n)$, to a mapping

$$F_{n,\ell}: R_{q,\ell}^b(n) \rightarrow \mathcal{P}\left((At^{-2\ell-4C} \cup At^{2\ell+4C}) \cap B_S(n + D')\right).$$

We prove below:

- (i) for each $g \in G$ with \mathcal{W} -itinerary $I = (\iota, \sigma)$ and for each $J \subseteq [\sigma^-, \sigma^+]_{\mathbb{Z}}$, the original element g can be recovered from $\dot{g}(J)$ and anyone of σ^+, σ^- ;
- (ii) for each $g \in G$ with \mathcal{W} -itinerary $I = (\iota, \sigma)$, the resulting elements are pairwise distinct: $\dot{g}(J) \neq \dot{g}(J')$ for $J, J' \subseteq [\sigma^-, \sigma^+]_{\mathbb{Z}}$ with $J \neq J'$;
- (iii) for every $h \in (At^{-2\ell-4C} \cup At^{2\ell+4C}) \cap B_S(n + D')$, where $\ell > q(n)$, there are at most $\ell + 1$ elements $g \in R_{q,\ell}^b(n)$ such that $h \in F_n(g)$.

From (ii) and (iii) above we conclude that

$$\left| (At^{-2\ell-4C} \cup At^{2\ell+4C}) \cap B_S(n + D') \right| \geq \frac{1}{\ell + 1} \binom{\ell + 1}{2} |R_{q,\ell}^b(n)| > \frac{q(n)}{2} |R_{q,\ell}^b(n)|.$$

Hence

$$|B_S(n + D')| > \frac{q(n)}{2} \sum_{\ell > q(n)} |R_{q,\ell}^b(n)| = \frac{q(n)}{2} |R_q^b(n)|,$$

which is the bound (3.5) we aimed for.

It remains to justify the statements (i), (ii) and (iii) above.

(i) Let $g \in G$ with itinerary $I = (\iota, \sigma)$, and let $J \subseteq [\sigma^-, \sigma^+]_{\mathbb{Z}}$. As in the discussion above suppose that $k^+ = k_{\mathcal{W},g}^+$ and $k^- = k_{\mathcal{W},g}^-$ satisfy $k^+ \leq k^-$; the other case $k^- < k^+$ can be dealt with similarly. It is not difficult to check that g can be recovered from $\dot{g}(J)$, if we are allowed to use one of the parameters σ^+, σ^- . Indeed,

from $-\rho(\dot{g}(J)) = 2(\sigma^+ - \sigma^-) + 4C$ we deduce that in such a case both, σ^+ and σ^- are available to us. Furthermore, Lemma 2.5(i) gives

$$\begin{aligned}\text{supp}(\dot{x}(J)) &\subseteq [\sigma^- - C + 1, \sigma^+ + C - 1]_{\mathbb{Z}}, \\ \text{supp}(\dot{y}(J)^{-1}) &\subseteq [-C + 1, \sigma^+ - \sigma^- + C - 1]_{\mathbb{Z}}, \\ \text{supp}(\dot{z}(J)) &\subseteq [-C + 1, -\sigma^- + C - 1]_{\mathbb{Z}},\end{aligned}$$

and thus

$$\begin{aligned}\text{supp}(\dot{g}(J)) &= \text{supp}(\dot{x}(J)) \cup (\text{supp}(\dot{y}(J)^{-1}) + \sigma^+ + 2C) \\ &\quad \cup (\text{supp}(\dot{z}(J)) + 2\sigma^+ - \sigma^- + 4C)\end{aligned}$$

allows us to recover $\dot{x}(J)$, $\dot{y}(J)$ and $\dot{z}(J)$ via (3.6) and

$$\begin{aligned}\dot{x}(J)|_i &= \begin{cases} \dot{g}(J)|_i & \text{for } i \in [\sigma^- - C, \sigma^+ + C]_{\mathbb{Z}}, \\ 1 & \text{for } i \in \mathbb{Z} \setminus [\sigma^- - C, \sigma^+ + C]_{\mathbb{Z}}, \end{cases} \\ (\dot{y}(J)^{-1})|_i &= \begin{cases} \dot{g}(J)|_{i+\sigma^++2C} & \text{for } i \in [-C, \sigma^+ - \sigma^- + C]_{\mathbb{Z}}, \\ 1 & \text{for } i \in \mathbb{Z} \setminus [-C, \sigma^+ - \sigma^- + C]_{\mathbb{Z}}, \end{cases} \\ \dot{z}(J)|_i &= \begin{cases} \dot{g}(J)|_{i+2\sigma^+-\sigma^-+4C} & \text{for } i \in [-C, -\sigma^- + C]_{\mathbb{Z}}, \\ 1 & \text{for } i \in \mathbb{Z} \setminus [-C, -\sigma^- + C]_{\mathbb{Z}}. \end{cases}\end{aligned}$$

Using (3.8), we recover $g = \dot{x}(J) \dot{y}(J) \dot{z}(J)$.

(ii) Let $g \in G$ with \mathcal{W} -itinerary $I = (\iota, \sigma)$. Again we suppose that $k^+ = k_{\mathcal{W},g}^+$ and $k^- = k_{\mathcal{W},g}^-$ satisfy $k^+ \leq k^-$; the other case $k^- < k^+$ can be dealt with similarly. Let $J, J' \subseteq [\sigma^-, \sigma^+]_{\mathbb{Z}}$ such that $\dot{g}(J) = \dot{g}(J')$. As explained above, we can not only recover g but even $\dot{x}(J) = \dot{x}(J')$, $\dot{y}(J) = \dot{y}(J')$ and $\dot{z}(J) = \dot{z}(J')$ from $\dot{g}(J) = \dot{g}(J')$ and σ^+ , say. Since $u \neq 1$ we deduce from (3.7) that $J = J'$.

(iii) First suppose that $h \in At^{2\ell+4C} \cap B_S(n + D')$, with $\ell > q(n)$, and suppose that $g \in R_{q,\ell}^b(n)$ such that $h = \dot{g}(J)$ for some $J \subseteq [\sigma_g^-, \sigma_g^+]_{\mathbb{Z}}$ with $|J| = 2$. Then $\sigma_g^+ \in [0, \ell]_{\mathbb{Z}}$ takes one of $\ell + 1$ values, and once σ^+ is fixed, there is a way of recovering g , by (i). For $h \in At^{-2\ell-4C} \cap B_S(n + D')$ the argument is similar. \square

4. PROOF OF THEOREM C

Throughout this section let G denote a finitely generated group of exponential word growth of the form $G = A \rtimes \langle t \rangle$, where

- (a) the subgroup $\langle t \rangle$ is infinite cyclic;
- (b) the subgroup $A = \langle \bigcup \{H^{t^i} \mid i \in \mathbb{Z}\} \rangle$ is generated by the $\langle t \rangle$ -conjugates of a finitely generated subgroup H ;
- (c) the $\langle t \rangle$ -conjugates of this group H commute elementwise: $[H^{t^i}, H^{t^j}] = 1$ for all $i, j \in \mathbb{Z}$ with $H^{t^i} \neq H^{t^j}$.

Suppose further that $S_0 = \{a_1, \dots, a_d\} \subseteq A$ is a finite symmetric generating set for H and that the exponential growth rates of H with respect to S_0 and of G with respect to $S = S_0 \cup \{t, t^{-1}\}$ satisfy

$$(4.1) \quad \lim_{n \rightarrow \infty} \sqrt[n]{|B_{H,S_0}(n)|} < \lim_{n \rightarrow \infty} \sqrt[n]{|B_{G,S}(n)|}.$$

This is essentially the setting of Theorem C; for technical reasons we prefer to work with symmetric generating sets. Our ultimate aim is to show that $\delta_S(A) = 0$.

Using the commutation rules recorded in (c), it is not difficult to see that every $g \in A$ admits S -expressions of minimal length that take the special form

$$(4.2) \quad g = t^{-\sigma^-} \cdot \prod_{i=\sigma^-}^{\sigma^+-1} (w_i(a_1, \dots, a_d) t^{-1}) \cdot w_{\sigma^+}(a_1, \dots, a_d) \cdot t^{\sigma^+},$$

$$(4.3) \quad g = t^{-\sigma^+} \cdot \prod_{j=\sigma^-}^{\sigma^+-1} (w_{\sigma^++\sigma^--j}(a_1, \dots, a_d) t) \cdot w_{\sigma^-}(a_1, \dots, a_d) \cdot t^{\sigma^-},$$

where the parameters $\sigma^-, \sigma^+ \in \mathbb{Z}$ satisfy $\sigma^- \leq \sigma^+$ and, for every $i \in [\sigma^-, \sigma^+]_{\mathbb{Z}}$, we have picked a suitable semigroup word $w_i = w_i(Y_1, \dots, Y_d)$ in d variables of length $l_{S_0}(w_i(a_1, \dots, a_d))$. The lengths of the expressions (4.2) and (4.3) are equal to

$$l_S(g) = |\sigma^-| + (\sigma^+ - \sigma^-) + |\sigma^+| + \sum_{i=\sigma^-}^{\sigma^+} l_{S_0}(w_i(a_1, \dots, a_d)).$$

For the following we fix, for each $g \in A$, expressions as described and we use subscripts to stress the dependency on g : we write σ_g^-, σ_g^+ and $w_{g,i}$ for $i \in [\sigma_g^-, \sigma_g^+]_{\mathbb{Z}}$, where necessary. The notation is meant to be reminiscent of the one introduced in Definition 2.2, but one needs to keep in mind that we are dealing with a larger class of groups now.

Lemma 4.1. *Let $q: \mathbb{N} \rightarrow \mathbb{R}_{>0}$ be a non-decreasing unbounded function such that $q \in o(n)$. For $n \in \mathbb{N}$ and the general set-up described above, we consider*

$$R_q(n) = \{g \in A \cap B_S(n) \mid -q(n) \leq \sigma_g^- \leq \sigma_g^+ \leq q(n)\}.$$

Then

$$\lim_{n \rightarrow \infty} \frac{|R_q(n)|}{|B_S(n)|} = 0.$$

Proof. For short we set $\mu = \lim_{n \rightarrow \infty} \sqrt[n]{|B_{H,S_0}(n)|}$ and $\lambda = \lim_{n \rightarrow \infty} \sqrt[n]{|B_{G,S}(n)|}$. According to (4.1) we find $\varepsilon \in \mathbb{R}_{>0}$ such that $(\mu + \varepsilon)/\lambda \leq 1 - \varepsilon$ and $M = M_\varepsilon \in \mathbb{N}$ such that

$$|B_{H,S_0}(n)| \leq M(\mu + \varepsilon)^n \quad \text{for all } n \in \mathbb{N}_0.$$

This allows us to bound the number of possibilities for the elements $w_{g,i}(a_1, \dots, a_d)$ in an S -expression of the form (4.2) for $g \in R_q(n)$ and, writing $\tilde{q}(n) = 2\lfloor q(n) \rfloor + 1$, we obtain

$$|R_q(n)| \leq \sum_{\substack{m_{-\lfloor q(n) \rfloor}, \dots, m_{\lfloor q(n) \rfloor} \in \mathbb{N}_0 \\ m_{-\lfloor q(n) \rfloor} + \dots + m_{\lfloor q(n) \rfloor} \leq n}} \prod_{i=-\lfloor q(n) \rfloor}^{\lfloor q(n) \rfloor} |B_{H,S_0}(m_i)| \leq \binom{n + \tilde{q}(n)}{\tilde{q}(n)} M^{\tilde{q}(n)} (\mu + \varepsilon)^n$$

and hence

$$(4.4) \quad \frac{|R_q(n)|}{|B_S(n)|} \leq \frac{|R_q(n)|}{\lambda^n} \leq \binom{n + \tilde{q}(n)}{\tilde{q}(n)} M^{\tilde{q}(n)} (1 - \varepsilon)^n \quad \text{for } n \in \mathbb{N}.$$

We notice that $q \in o(n)$ implies $\tilde{q} \in o(n)$. Thus Lemma 2.1 implies that $\binom{n + \tilde{q}(n)}{\tilde{q}(n)} M^{\tilde{q}(n)}$ grows sub-exponentially, and the term on the right-hand side of (4.4) tends to 0 as n tends to infinity. \square

Proof of Theorem C. We continue to work in the notational set-up introduced above. In addition we fix a non-decreasing unbounded function $q: \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$ such that $q \in o(n)$; e.g., $q(n) = n/(\log(n) + 1)$ works nicely. As in the proof of Theorem B, we make use of a decomposition

$$A \cap B_S(n) = R_q(n) \cup R_q^b(n), \quad \text{for } n \in \mathbb{N},$$

where $R_q(n)$ is defined as in Lemma 4.1 and $R_q^b(n)$ denotes the corresponding complement.

In view of Lemma 4.1 it suffices to show that

$$(4.5) \quad \frac{|R_q^b(n)|}{|B_S(n)|} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Indeed, for any $g \in R_q^b(n)$, with chosen minimal S -expressions (4.2) and (4.3), we have $\sigma^- = \sigma_g^- < -q(n)$ or $\sigma^+ = \sigma_g^+ > q(n)$, hence

$$\left\{gt^{-q(n)}, gt^{q(n)}\right\} \cap B_S(n - q(n)) \neq \emptyset.$$

As each of the right translation maps $g \mapsto gt^{-q(n)}$ and $g \mapsto gt^{q(n)}$ is injective, we conclude that

$$|R_q^b(n)| \leq 2|B_S(n - q(n))|$$

and thus, for any $m \in \mathbb{N}$ and all $n \in \mathbb{N}$ with $q(n) \geq m$,

$$\frac{|R_q^b(n)|}{|B_S(n)|} \leq \frac{2|B_S(n - q(n))|}{|B_S(n)|} \leq \frac{2|B_S(n - m)|}{|B_S(n)|} \rightarrow \frac{2}{\lambda^m} \quad \text{as } n \rightarrow \infty.$$

Since $\lambda > 1$ and q is unbounded, we conclude that (4.5) holds. \square

REFERENCES

- [1] Antolín, Y., Martino, A. and Ventura, E.: Degree of commutativity of infinite groups, *Proc. Amer. Math. Soc.* **145**(2) (2017), 479–485.
- [2] Burillo, J. and Ventura, E.: Counting primitive elements in free groups, *Geom. Dedicata* **93** (2002), 143–162.
- [3] Cox, C. G.: The degree of commutativity and lamplighter groups, *Internat. J. Algebra Comput.* **28**(7) (2018), 1163–1173.
- [4] Erdős, P. and Turán, P.: On some problems in statistical group-theory IV, *Acta Math. Acad. Sci. Hung.* **19** (1968), 413–435.
- [5] Guralnick, R. M. and Robinson, G. R.: On the commuting probability in finite groups, *J. Algebra* **300** (2006), 509–528.
- [6] Gustafson, W. H.: What is the probability that two group elements commute?, *Amer. Math. Monthly* **80**(9) (1973), 1031–1034.
- [7] Neumann, P. M.: Two combinatorial problems in group theory, *Bull. London Math. Soc.* **21** (1989), 456–458.
- [8] Rusin, D. J.: What is the probability that two elements of a finite group commute?, *Pacific J. Math.* **82**(1) (1979), 237–247.
- [9] Shalev, A.: Probabilistically nilpotent groups, *Proc. Amer. Math. Soc.* **146** (2018), 1529–1536.
- [10] Tointon, M. C. H.: Commuting probabilities of infinite groups, *J. London Math. Soc.* **101**(3) (2020), 1280–1297.
- [11] Valiunas, M.: Rational growth and degree of commutativity of graph products, *J. Algebra* **522** (2019), 309–331.

IKER DE LAS HERAS: MATHEMATISCHES INSTITUT, HEINRICH-HEINE-UNIVERSITÄT, 40225 DÜSSELDORF, GERMANY; DEPARTMENT OF MATHEMATICS, UNIVERSITY OF THE BASQUE COUNTRY UPV/EHU, 48940 LEIOA, SPAIN

Email address: `iker.delasheras@hhu.de`

BENJAMIN KLOPSCH: MATHEMATISCHES INSTITUT, HEINRICH-HEINE-UNIVERSITÄT, 40225 DÜSSELDORF, GERMANY

Email address: `klopsch@math.uni-duesseldorf.de`

ANDONI ZOZAYA: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF THE BASQUE COUNTRY UPV/EHU, 48940 LEIOA, SPAIN

Email address: `andoni.zozaya@ehu.eus`